

REPRESENTATION FOR BOUNDED LINEAR OPERATOR ON HILBERT SPACES

LVLIN LUO *

ABSTRACT. In this paper we construct some C^* -algebra induced by polar decomposition $T = U|T|$. We get that T is unitary equivalent to $\sqrt{|\eta|}M_{z\phi}$ on $\mathcal{L}^2(\sigma(|T|), \mu_{|T|})$, where $\phi \in \mathcal{L}^\infty(\sigma(|T|), \mu_{|T|})$ and $\eta \in \mathcal{L}^1(\sigma(|T|), \mu_{|T|})$. Also, we get that T is normal if and only if T is unitary equivalent to $M_{z\phi}$ on $\mathcal{L}^2(\sigma(|T|), \mu_{|T|})$, and if and only if $T \in \mathcal{A}'(|T|)$, where $\phi \in \mathcal{L}^\infty(\sigma(|T|), \mu_{|T|})$ and $\mathcal{A}'(|T|)$ is the commutant of $\mathcal{A}(|T|)$.

1. INTRODUCTION

Linear Dynamics is a rapidly evolving branch of functional analysis, which was probably born in 1982 with the Toronto Ph. D. thesis of C. Kitai [1]. It has become rather popular because of the efforts of many mathematicians survey [5, 11, 15, 4, 16]. Today, interdiscipline becomes a common phenomenon in mathematics. And there is new objects about C^* -algebras and dynamics, such as [24] in 2014 and [25] in 2015. In [22], Gelfand-Naimark theorem [10] is called “the fundamental theorem of C^* -algebras”. Such that Ichiro Fujimoto said that this theorem eventually opened the gate to the subject of C^* -algebras [9]. Hence there are various attempts to generalize this theorem [12, 19, 3, 8, 21].

For invertible dynamics, the relationship between (X, f) and (X, f^{-1}) is raised by Stockman as an open question [2]. And [18, 7] and [17] give counter example with some chaos for this question in noncompact spaces and compact spaces, respectively. For invertible bounded linear operator on Hilbert spaces, the dynamics relationship between T and T^{*-1} is also interested. Any way, counter example is trivial for some chaos between T and T^{*-1} . But there is no generally way to this research. In fact, the C^* -algebra $\mathcal{A}(T)$ generated by T is not obvious effect for that. So we get another way, that is representation theory for T by C^* -algebra. And which is one step to get some dynamics relationship between T and T^{*-1} .

2. PRELIMINARIES

The Polar Decomposition theorem [14]P15 on Hilbert space is a useful theorem, especially for invertible bounded linear operator. And we give some properties of C^* -algebra generated by normal operator.

Date: Received: xxxxxx; Revised: yyyyyy; Accepted: zzzzzz.

* Corresponding author.

2010 *Mathematics Subject Classification.* Primary 37B99, 46L05, 47A65, 47A67; Secondary

Key words and phrases. polar decomposition; spectrum; C^* algebra; unitary equivalent.

Let $\mathbb{H}$ be a separable Hilbert space over $\mathbb{C}$ and let X be a compact subset of $\mathbb{C}$. Let $\mathcal{C}(X)$ denote the linear space of all continuous functions on the compact space X and let $\mathcal{P}(x)$ denote the set of all polynomials on X . Let T be a bounded linear operator on $\mathbb{H}$ and $\sigma(T)$ is its spectrum. By the Polar Decomposition theorem [14]P15 we get $T = U|T|$ and $|T|^2 = T^*T$. When T is invertible, U is a unitary operator. Let $\mathcal{A}(|T|)$ denote the C^* -algebra generated by $|T|$ and 1.

Lemma 2.1. *Let $0 \notin X$ be a compact subset of $\mathbb{C}$. If $\mathcal{P}(x)$ is dense in $\mathcal{C}(X)$, then $\mathcal{P}(\frac{1}{x})$ is also dense in $\mathcal{C}(X)$.*

Proof. By the property of polynomials we know that $\mathcal{P}(\frac{1}{x})$ is a algebraic closed subalgebra of $\mathcal{C}(X)$ and we get:

- (1) $1 \in \mathcal{P}(\frac{1}{x})$;
- (2) $\mathcal{P}(\frac{1}{x})$ separate the points of X ;
- (3) If $p(\frac{1}{x}) \in \mathcal{P}(x)$, then $\bar{p}(\frac{1}{x}) \in \mathcal{P}(x)$.

By the Stone-Weierstrass theorem [13]P145 we get the conclusion. $\square$

Lemma 2.2. *Let $X \subseteq \mathbb{R}_+$. If $\mathcal{P}(|x|)$ is dense in $\mathcal{C}(X)$, then $\mathcal{P}(|x|^2)$ is also dense in $\mathcal{C}(X)$.*

Proof. For $X \subseteq \mathbb{R}_+$, $x \neq y \iff x^2 \neq y^2$. By Lemma 2.1 we get the conclusion. $\square$

3. DECOMPOSITION AND ISOMORPHIC

By the GNS construction [13]P250 for the C^* -algebra $\mathcal{A}(|T|)$, we have the following decomposition.

Lemma 3.1. *Let T be an invertible bounded linear operator on the separable Hilbert space $\mathbb{H}$ over $\mathbb{C}$, $\mathcal{A}(|T|)$ is the complex C^* -algebra generated by $|T|$ and 1. There is a sequence of nonzero $\mathcal{A}(|T|)$ -invariant subspace. $\mathbb{H}_1, \mathbb{H}_2, \dots$ such that:*

- (1) $\mathbb{H} = \mathbb{H}_1 \oplus \mathbb{H}_2 \oplus \dots$;
- (2) For every $\mathbb{H}_i$, there is a $\mathcal{A}(|T|)$ -cyclic vector ξ^i such that $\mathbb{H}_i = \overline{\mathcal{A}(|T|)\xi^i}$ and $|T|\mathbb{H}_i = \mathbb{H}_i = |T|^{-1}\mathbb{H}_i$.

Proof. The decomposition of (1) is obvious by [23]P54. Hence $|T|\mathbb{H}_i \subseteq \mathbb{H}_i$, that is $\mathbb{H}_i \subseteq |T|^{-1}\mathbb{H}_i$. By Lemma 2.1 we get $|T|^{-1}\mathbb{H}_i \subseteq \mathbb{H}_i$. Therefore $|T|\mathbb{H}_i = \mathbb{H}_i = |T|^{-1}\mathbb{H}_i$. $\square$

For $\forall n \in \mathbb{N}$, T^n is invertible when T is invertible. With the Polar Decomposition theorem [14]P15 $T^n = U_n|T^n|$, where U_n is unitary operator and $|T^n|^2 = T^{*n}T^n$, we get the following conclusion.

Lemma 3.2. *Let T be an invertible bounded linear operator on the separable Hilbert space $\mathbb{H}$ over $\mathbb{C}$, let $\mathcal{A}(|T^k|)$ be the complex C^* -algebra generated by $|T^k|$ and 1 and let $\mathbb{H}_i^{[T^k]} = \overline{\mathcal{A}(|T^k|)\xi_k^i}$ be a sequence of non-zero $\mathcal{A}(|T^k|)$ -invariant subspace, there is a decomposition $\mathbb{H} = \bigoplus_i \mathbb{H}_i^{[T^k]}$, $\xi_k^i \in \mathbb{H}$, $i, k \in \mathbb{N}$. Given a proper permutation of $\mathbb{H}_i^{[T^k]}$ and $\mathbb{H}_j^{[T^{(k+1)}]}$, we get $T^*\mathbb{H}_i^{[T^k]} = \mathbb{H}_i^{[T^{(k+1)}]}$ and $T^{-1}\mathbb{H}_i^{[T^k]} = \mathbb{H}_i^{[T^{(k+1)}]}$.*

Proof. Using Lemma 2.2, it is enough to prove the conclusion on $\overline{\mathcal{P}(|T^k|^2)\xi_k^i}$. For any given $\xi_k^i \in \mathbb{H} = \bigoplus_j \mathbb{H}_j^{|T^{(k+1)}|}$, there is a unique $j \in \mathbb{N}$ such that $\xi_k^i \in \mathbb{H}_j^{|T^{(k+1)}|}$.

(1) Because T is invertible, for any given ξ_k^i , there is a unique $\eta_i \in \mathbb{H}_s^{|T^{(k+1)}|}$ such that $\eta_i = T^{-1}\xi_k^i$. For $\forall p \in \mathcal{P}(|x|^2)$, we get $p(|T^{(k+1)}|^2)\eta_i = T^*p(|T^k|^2)\xi_k^i$. Hence

$$\mathbb{H}_s^{|T^{(k+1)}|} = \overline{\mathcal{P}(|T^{(k+1)}|^2)\xi_{k+1}^s} \supseteq T^*\overline{\mathcal{P}(|T^k|^2)\xi_k^i} = T^*\mathbb{H}_i^{|T^k|}.$$

(2) Similarly, for any given ξ_{k+1}^s , there is a unique $\eta_r \in \mathbb{H}_r^{|T^k|}$ such that $\eta_r = T\xi_{k+1}^s$. For $\forall p \in \mathcal{P}(|x|^2)$, we get

$$p(|T^k|^2)\eta_r = p(|T^k|^2)T\xi_{k+1}^s = T^{*-1}p(|T^{(k+1)}|^2)\xi_{k+1}^s.$$

Hence

$$\mathbb{H}_r^{|T^k|} = \overline{\mathcal{P}(|T^k|^2)\xi_k^r} \supseteq T^{*-1}\overline{\mathcal{P}(|T^{(k+1)}|^2)\xi_{k+1}^s} = T^{*-1}\mathbb{H}_s^{|T^{(k+1)}|}.$$

Let $i = r$, by (1)(2) we get

$$T^{*-1}\mathbb{H}_s^{|T^{(k+1)}|} \subseteq \mathbb{H}_i^{|T^k|} \subseteq T^{*-1}\mathbb{H}_j^{|T^{(k+1)}|}.$$

Fixed the order of $\mathbb{H}_i^{|T^k|}$, by a proper permutation of $\mathbb{H}_j^{|T^{(k+1)}|}$ we get

$$T^*\mathbb{H}_i^{|T^k|} = \mathbb{H}_i^{|T^{(k+1)}|}.$$

By Lemma 2.1 and T is invertible, there are the following next conclusion.

(3) For any given ξ_k^i , there is a unique $\eta_i \in \mathbb{H}_i^{|T^{(k+1)}|}$ such that $\eta_i = T^*\xi_k^i$. For $\forall p \in \mathcal{P}(|x|^{-2})$, we get $p(|T^{(k+1)}|^{-2})\eta_i = T^{-1}p(|T^k|^{-2})\xi_k^i$. Hence

$$\mathbb{H}_i^{|T^{(k+1)}|} = \overline{\mathcal{P}(|T^{(k+1)}|^{-2})\xi_{k+1}^i} \supseteq T^{-1}\overline{\mathcal{P}(|T^k|^{-2})\xi_k^i} = T^{-1}\mathbb{H}_i^{|T^k|}.$$

(4) For any given ξ_{k+1}^i , there is a unique $\eta_i \in \mathbb{H}_i^{|T^k|}$ such that $\eta_i = T^{*-1}\xi_{k+1}^i$. For $\forall p \in \mathcal{P}(|x|^{-2})$, we get

$$T^{-1}p(|T^k|^{-2})\eta_i = T^{-1}p(|T^k|^{-2})T^{*-1}\xi_{k+1}^i = p(|T^{(k+1)}|^{-2})\xi_{k+1}^i.$$

Hence

$$T^{-1}\mathbb{H}_i^{|T^k|} = T^{-1}\overline{\mathcal{P}(|T^k|^{-2})\xi_k^i} \supseteq \overline{\mathcal{P}(|T^{(k+1)}|^{-2})\xi_{k+1}^i} = \mathbb{H}_i^{|T^{(k+1)}|}.$$

By (3)(4) we get

$$T^{-1}\mathbb{H}_i^{|T^k|} \subseteq \mathbb{H}_i^{|T^{(k+1)}|} \subseteq T^{-1}\mathbb{H}_i^{|T^k|}.$$

that is, $T^{-1}\mathbb{H}_i^{|T^k|} = \mathbb{H}_i^{|T^{(k+1)}|}$. □

Let $\xi \in \mathbb{H}$ is a $\mathcal{A}(|T|)$ -cyclic vector such that $\mathcal{A}(|T|)\xi$ is dense in $\mathbb{H}$. Because of the spectral $\sigma(|T|) \neq \emptyset$, on $\mathcal{C}(\sigma(|T|))$ define the non-zero linear functional

$$\rho_{|T|} : \rho_{|T|}(f) = \langle f(|T|)\xi, \xi \rangle, \forall f \in \mathcal{C}(\sigma(|T|)).$$

Then $\rho_{|T|}$ is a positive linear functional, by [23]P54 and the Riesz-Markov theorem, on $\mathcal{C}(\sigma(|T|))$ we get that there exists an unique finite positive Borel measure $\mu_{|T|}$ such that

$$\int_{\sigma(|T|)} f(z) d\mu_{|T|}(z) = \langle f(|T|)\xi, \xi \rangle, \quad \forall f \in \mathcal{C}(\sigma(|T|)).$$

Theorem 3.3. *Let T be an invertible bounded linear operator on the separable Hilbert space $\mathbb{H}$ over $\mathbb{C}$, there is $\xi \in \mathbb{H}$ such that $\overline{\mathcal{A}(|T|)}\xi = \mathbb{H}$. For any given $n \in \mathbb{N}$, let $\mathcal{A}(|T^n|)$ be the complex C^* -algebra generated by $|T^n|$ and 1 and let ξ_n be a $\mathcal{A}(|T^n|)$ -cyclic vector such that $\overline{\mathcal{A}(|T^n|)}\xi_n = \mathbb{H}$. Then:*

(1) *For any given ξ_n , there is an unique positive linear functional*

$$\int_{\sigma(|T^n|)} f(z) d\mu_{|T^n|}(z) = \langle f(|T^n|)\xi_n, \xi_n \rangle, \quad \forall f \in \mathcal{L}^2(\sigma(|T^n|)).$$

(2) *For any given ξ_n , there is an unique finite positive complete Borel measure $\mu_{|T^n|}$ such that $\mathcal{L}^2(\sigma(|T^n|))$ is isomorphic to $\mathbb{H}$.*

Proof. Because T is invertible, by Lemma 3.2 we get that if there is a $\mathcal{A}(|T|)$ -cyclic vector ξ , then there is a $\mathcal{A}(|T^n|)$ -cyclic vector ξ_n .

(1) For any given ξ_n , define the linear functional

$$\rho_{|T^n|}(f) = \langle f(|T^n|)\xi_n, \xi_n \rangle,$$

by [23]P54 and the Riesz-Markov theorem we get that on $\mathcal{C}(\sigma(|T^n|))$ there is an unique finite positive Borel measure $\mu_{|T^n|}$ such that

$$\int_{\sigma(|T^n|)} f(z) d\mu_{|T^n|}(z) = \langle f(|T^n|)\xi_n, \xi_n \rangle, \quad \forall f \in \mathcal{C}(\sigma(|T^n|)).$$

Moreover we can complete the Borel measure $\mu_{|T^n|}$ on $\sigma(|T^n|)$, also using $\mu_{|T^n|}$ to denote the complete Borel measure, By [20] we know that the complete Borel measure is uniquely.

For $\forall f \in \mathcal{L}^2(\sigma(|T^n|))$, because of

$$\rho_{|T^n|}(|f|^2) = \rho_{|T^n|}(\bar{f}f) = \langle f(|T^n|)^*f(|T^n|)\xi_n, \xi_n \rangle = \|f(|T^n|)\xi_n\|^2 \geq 0.$$

we get that $\rho_{|T^n|}$ is a positive linear functional, that is (1).

(2) we know that $\mathcal{C}(\sigma(|T^n|))$ is a subspace of $\mathcal{L}^2(\sigma(|T^n|))$ such that $\mathcal{C}(\sigma(|T^n|))$ is dense in $\mathcal{L}^2(\sigma(|T^n|))$. For any $f, g \in \mathcal{C}(\sigma(|T^n|))$ we get

$$\begin{aligned} & \langle f(|T^n|)\xi_n, g(|T^n|)\xi_n \rangle_{\mathbb{H}} \\ &= \langle g(|T^n|)^*f(|T^n|)\xi_n, \xi_n \rangle \\ &= \rho_{|T^n|}(\bar{g}f) \\ &= \int_{\sigma(|T^n|)} f(z)\bar{g}(z) d\mu_{|T^n|}(z) \\ &= \langle f, g \rangle_{\mathcal{L}^2(\sigma(|T^n|))}. \end{aligned}$$

Therefor $U_0 : \mathcal{C}(\sigma(|T^n|)) \rightarrow \mathbb{H}, f(z) \rightarrow f(|T^n|)\xi_n$ is a surjection isometry from $\mathcal{C}(\sigma(|T^n|))$ to $\mathcal{A}(|T^n|)\xi_n$, also $\mathcal{C}(\sigma(|T^n|))$ and $\mathcal{A}(|T^n|)\xi_n$ is a dense subspace of $\mathcal{L}^2(\sigma(|T^n|))$ and $\mathbb{H}$, respectively. Because of U_0 a closable operator, it's closed extension $U : \mathcal{L}^2(\sigma(|T^n|)) \rightarrow \mathbb{H}, f(z) \rightarrow f(|T^n|)\xi_n$ is a unitary operator. Hence for any given ξ_n , U is the unique unitary operator induced by the unique finite positive complete Borel measure $\mu_{|T^n|}$ such that $\mathcal{L}^2(\sigma(|T^n|))$ is isomorphic to $\mathbb{H}$. $\square$

4. UNITARY OPERATOR AND REPRESENTATION

By the Polar Decomposition theorem $T = U|T|$ [14]P15, we get $U^*T^*TU = TT^*$ and $U^*|T|^{-2}U = |T^{-2}|$. In fact, when T is invertible, we can choose an specially unitary operator such that $|T|^{-1}$ and $|T^{-1}|$ are unitary equivalent. We have the following unitary equivalent by Theorem 3.3.

Theorem 4.1. *Let T be an invertible bounded linear operator on the separable Hilbert space $\mathbb{H}$ over $\mathbb{C}$ and let $\mathcal{A}(|T|)$ be the complex C^* -algebra generated by $|T|$ and 1. There is $\sigma(|T|^{-1}) = \sigma(|T^{-1}|)$ and we get that $|T|^{-1}$ and $|T^{-1}|$ are unitary equivalent by the unitary operator $F_{xx^*}^{\mathbb{H}}$, more over the unitary operator $F_{xx^*}^{\mathbb{H}}$ is induced by an almost everywhere non-zero function $\sqrt{|\phi_{|T|}|}$, where $\sqrt{|\phi_{|T|}|} \in \mathcal{L}^2(\sigma(|T|), \mu_{|T|})$. That is, $d\mu_{|T^{-1}|} = |\phi_{|T|}|d\mu_{|T|^{-1}}$.*

Proof. By Lemma 3.1, lose no generally, let $\xi_{|T|}$ is a $\mathcal{A}(|T|)$ -cyclic vector such that $\mathbb{H} = \overline{\mathcal{A}(|T|)\xi_{|T|}}$.

(1) Define the function

$$F_{x^{-1}} : \mathcal{P}(x) \rightarrow \mathcal{P}(x^{-1}), F_{x^{-1}}(f(x)) = f(x^{-1}),$$

it is easy to find that $F_{x^{-1}}$ is linear. Because of

$$\int_{\sigma(|T|)} f(z^{-1})d\mu_{|T|}(z) = \langle f(|T|^{-1})\xi, \xi \rangle = \int_{\sigma(|T|^{-1})} f(z)d\mu_{|T|^{-1}}(z).$$

We get $d\mu_{|T|^{-1}}(z) = |z|^2d\mu_{|T|}(z)$. Hence

$$\begin{aligned}
& \|F_{x^{-1}}(f(x))\|_{\mathcal{L}^2(\sigma(|T|^{-1}), \mu_{|T|^{-1}})}^2 \\
&= \int_{\sigma(|T|^{-1})} F_{x^{-1}}(f(x)) \bar{F}_{x^{-1}}(f(x)) d\mu_{|T|^{-1}}(x) \\
&= \int_{\sigma(|T|^{-1})} f(x^{-1}) \bar{f}(x^{-1}) d\mu_{|T|^{-1}}(x) \\
&= \int_{\sigma(|T|)} |z|^2 f(z) \bar{f}(z) d\mu_{|T|}(z) \\
&\leq \sup \sigma(|T|)^2 \int_{\sigma(|T|)} f(z) \bar{f}(z) d\mu_{|T|}(z) \\
&\leq \sup \sigma(|T|)^2 \|f(z)\|_{\mathcal{L}^2(\sigma(|T|), \mu_{|T|})}^2.
\end{aligned}$$

Therefor $\|F_{x^{-1}}\| \leq \sup \sigma(|T|)$, by the Banach Inverse Mapping theorem [13]P91 we get that $F_{x^{-1}}$ is an invertible bounded linear operator from $\mathcal{L}^2(\sigma(|T|), \mu_{|T|})$ to $\mathcal{L}^2(\sigma(|T|^{-1}), \mu_{|T|^{-1}})$.

Define the operator

$$F_{x^{-1}}^{\mathbb{H}} : \mathcal{A}(|T|)\xi \rightarrow \mathcal{A}(|T|^{-1})\xi, F(f(|T|)\xi) = f(|T|^{-1})\xi.$$

By Lemma 2.1 and [23]P55, we get that $F_{x^{-1}}^{\mathbb{H}}$ is a bounded linear operator on the Hilbert space $\overline{\mathcal{A}(|T|)\xi} = \mathbb{H}$, $\|F_{x^{-1}}^{\mathbb{H}}\| \leq \sup \sigma(|T|)$. Moreover we get

$$\begin{array}{ccc}
\mathbb{H} & \xrightarrow{|T|} & \mathbb{H} \\
F_{x^{-1}}^{\mathbb{H}} \downarrow & & \downarrow F_{x^{-1}}^{\mathbb{H}} \\
\mathbb{H} & \xrightarrow{|T|^{-1}} & \mathbb{H}
\end{array}$$

(2) Define the function

$$F_{xx^*} : \mathcal{P}(xx^*) \rightarrow \mathcal{P}(x^*x), F_{xx^*}(f(x, x^*)) = f(x^*x).$$

By [6] we get that F_{xx^*} is a linear algebraic isomorphism. For any $x \in \sigma(|T^{-1}|)$, by Lemma 2.2 we get that $\mathcal{P}(|x|^2)$ is dense in $\mathcal{C}(|x|)$ and $\mathcal{C}(|x|)$ is dense in $\mathcal{L}^2(\sigma(|T^{-1}|), \mu_{|T^{-1}|})$. Hence $\mathcal{P}(|x|^2)$ is dense in $\mathcal{L}^2(\sigma(|T^{-1}|), \mu_{|T^{-1}|})$.

With a similarly discussion, for any $y \in \sigma(|T|^{-1})$, we get that $\mathcal{P}(|y|^2)$ is dense in $\mathcal{L}^2(\sigma(|T|^{-1}), \mu_{|T|^{-1}})$.

So F_{xx^*} is an invertible bounded linear operator from $\mathcal{L}^2(\sigma(|T|^{-1}), \mu_{|T|^{-1}})$ to $\mathcal{L}^2(\sigma(|T|), \mu_{|T|})$. Therefor $F_{xx^*} \circ F_{x^{-1}}$ is an invertible bounded linear operator from $\mathcal{L}^2(\sigma(|T|), \mu_{|T|})$ to $\mathcal{L}^2(\sigma(|T^{-1}|), \mu_{|T^{-1}|})$.

Because of

$$\lambda \in \sigma(T^*T) \iff \frac{1}{\lambda} \in \sigma(T^{*-1}T^{-1}),$$

we get

$$\lambda \in \sigma(|T|) \iff \frac{1}{\lambda} \in \sigma(|T^{-1}|),$$

that is,

$$\sigma(|T^{-1}|) = \sigma(|T|^{-1}).$$

For any $p_n \in \mathcal{P}(\sigma(|T|^{-1})) \subseteq \mathcal{A}(\sigma(|T|^{-1}))$, because of $T^{*-1}p_n(|T|^{-1}) = p_n(|T^{-1}|)T^{*-1}$, by [14]P60 we get that $\mathcal{P}(|T|^{-1})$ and $\mathcal{P}(|T^{-1}|)$ are unitary equivalent. Hence there is an unitary operator $U \in \mathcal{B}(\mathbb{H})$ such that $U\mathcal{P}(|T|^{-1}) = \mathcal{P}(|T^{-1}|)U$, that is, $U\mathcal{A}(|T|^{-1}) = \mathcal{A}(|T^{-1}|)U$ and $U\overline{\mathcal{A}(|T|^{-1})}\xi_{|T|} = \overline{\mathcal{A}(|T^{-1}|)}U\xi_{|T|}$. Let $\xi_{|T^{-1}|} = U\xi_{|T|}$, then $\xi_{|T^{-1}|}$ is a $\mathcal{A}(|T^{-1}|)$ -cyclic vector and $\overline{\mathcal{A}(|T^{-1}|)}\xi_{|T^{-1}|} = \mathbb{H}$. Because of

$$\begin{aligned} & \int_{\sigma(|T|^{-1})} f(z) d\mu_{|T|^{-1}}(z) \\ &= \langle f(|T|^{-1})\xi_{|T|}, \xi_{|T|} \rangle \\ &= \int_{\sigma(|T|)} f\left(\frac{1}{z}\right) d\mu_{|T|}(z) \end{aligned}$$

and

$$\begin{aligned} & \int_{\sigma(|T^{-1}|)} f(z) d\mu_{|T^{-1}|}(z) \\ &= \langle f(|T^{-1}|)\xi_{|T^{-1}|}, \xi_{|T^{-1}|} \rangle, \end{aligned}$$

also F_{xx^*} is a linear algebraic isomorphic, we get $[d\mu_{|T^{-1}|}] = [d\mu_{|T|^{-1}}]$, that is, $d\mu_{|T^{-1}|}$ and $d\mu_{|T|^{-1}}$ are mutually absolutely continuous, by [13]IX.3.6Theorem and (1) we get

$$d\mu_{|T^{-1}|} = |\phi_{|T|}\left(\frac{1}{z}\right)| d\mu_{|T|^{-1}} = |z|^2 |\phi_{|T|}(z)| d\mu_{|T|},$$

where $|\phi_{|T|}(z)| \neq 0, a.e.$ and $\phi_{|T|}(z) \in \mathcal{L}^1(\sigma(|T|), \mu_{|T|})$. So we get

$$\begin{aligned}
& \|F_{xx^*} \circ F_{x^{-1}}(f(x))\|_{\mathcal{L}^2(\sigma(|T^{-1}|), \mu_{|T^{-1}|})}^2 \\
&= \int_{\sigma(|T^{-1}|)} F_{xx^*} \circ F_{x^{-1}}(f(x)) \overline{F_{xx^*} \circ F_{x^{-1}}(f(x))} d\mu_{|T^{-1}|}(x) \\
&= \int_{\sigma(|T^{-1}|)} f(x^{-1}) \bar{f}(x^{-1}) d\mu_{|T^{-1}|}(x) \\
&= \int_{\sigma(|T|^{-1})} |f(x^{-1}) \bar{f}(x^{-1})| \phi_{|T|}(\frac{1}{x}) d\mu_{|T|^{-1}}(x) \\
&= \int_{\sigma(|T|^{-1})} F_{x^{-1}}(f(x)) \sqrt{|\phi_{|T|}(x)|} F_{x^{-1}}(\bar{f}(x)) \sqrt{|\phi_{|T|}(x)|} d\mu_{|T|^{-1}}(x) \\
&= \|F_{x^{-1}}(f(x)) \sqrt{|\phi_{|T|}(\frac{1}{x})|}\|_{\mathcal{L}^2(\sigma(|T|^{-1}), \mu_{|T|^{-1}})}^2
\end{aligned}$$

Hence F_{xx^*} is a unitary operator that is induced by $Uf(\frac{1}{x}) = f(\frac{1}{x})\sqrt{|\phi_{|T|}(\frac{1}{x})|}$. Define

$$F_{xx^*}^{\mathbb{H}} : \mathcal{A}(|T|^{-2})\xi_{|T|} \rightarrow \mathcal{A}(|T^{-2}|)\xi_{|T^{-1}|}, F_{xx^*}^{\mathbb{H}}(f(|T|^{-2})\xi_{|T|}) = f(|T^{-2}|)\xi_{|T^{-1}|}.$$

Therefor $F_{xx^*}^{\mathbb{H}}$ is a unitary operator from $\overline{\mathcal{A}(|T|^{-1})\xi_{|T|}}$ to $\overline{\mathcal{A}(|T^{-1}|)\xi_{|T^{-1}|}}$. By Lemma 3.1 and [6] we get

$$\overline{\mathcal{A}(|T|^{-1})\xi_{|T|}} = \mathbb{H} = \overline{\mathcal{A}(|T^{-1}|)\xi_{|T^{-1}|}}.$$

That is, $F_{xx^*}^{\mathbb{H}}$ is a unitary operator and we get

$$\begin{array}{ccc}
\mathbb{H} & \xrightarrow{|T|^{-1}} & \mathbb{H} \\
F_{xx^*}^{\mathbb{H}} \downarrow & & \downarrow F_{xx^*}^{\mathbb{H}} \\
\mathbb{H} & \xrightarrow{|T^{-1}|} & \mathbb{H}
\end{array}$$

So $|T|^{-1}$ and $|T^{-1}|$ are unitary equivalent by $F_{xx^*}^{\mathbb{H}}$, the unitary operator $F_{xx^*}^{\mathbb{H}}$ is induced by the function $\sqrt{|\phi_{|T|}(\frac{1}{x})|}, \sqrt{|\phi_{|T|}(\frac{1}{x})|} \in \mathcal{L}^2(\sigma(|T|^{-1}), \mu_{|T|^{-1}})$. $\square$

Corollary 4.2. *Let T be an invertible bounded linear operator on the separable Hilbert space $\mathbb{H}$ over $\mathbb{C}$ and let $\mathcal{A}(|T|)$ be the complex C^* -algebra generated by $|T|$ and 1. There is $\sigma(|T|) = \sigma(|T^*|)$ and we get that $|T|$ and $|T^*|$ are unitary equivalent by the unitary operator $F_{xx^*}^{\mathbb{H}}$, more over the unitary operator $F_{xx^*}^{\mathbb{H}}$ is induced by an almost everywhere non-zero function $\sqrt{|\phi_{|T|}|}$, where $\sqrt{|\phi_{|T|}|} \in \mathcal{L}^2(\sigma(|T|), \mu_{|T|})$. That is, $d\mu_{|T^*|} = |\phi_{|T|}|d\mu_{|T|}$.*

Theorem 4.3. *Let T be an invertible bounded linear operator on the separable Hilbert space $\mathbb{H}$ over $\mathbb{C}$, $T = U|T|$ is the Polar Decomposition, and M_z is the unitary equivalent representation of $|T|$ on $\mathcal{L}^2(\sigma(|T|), \mu_{|T|})$, where define*

M_z is $M_z f = z f$ on $\mathcal{L}^2(\sigma(|T|), \mu_{|T|})$. There is $\phi \in \mathcal{L}^1(\sigma(|T|), \mu_{|T|})$ and $\psi \in \mathcal{L}^\infty(\sigma(|T|), \mu_{|T|})$ such that $\sqrt{|\phi|} M_{z\psi}$ is the unitary equivalent representation of T on $\mathcal{L}^2(\sigma(|T|), \mu_{|T|})$, and $\sqrt{|\phi|} M_\psi$ is the unitary equivalent representation of the unitary operator U .

Proof. By the Polar Decomposition $T = U|T|$, we get that $T^*T = |T|^2$ and $TT^* = U|T|^2U^*$. By Corollary 4.2 we get $TT^* = F_{xx^*}^{\mathbb{H}}|T|^2(F_{xx^*}^{\mathbb{H}})^*$. With the fact that $\{M_\phi : \phi \in \mathcal{L}^\infty(\sigma(|T|), \mu_{|T|})\}$ is a maximal abelian von Neumann algebra in $\mathcal{B}(\mathcal{L}^2(\sigma(|T|), \mu_{|T|}))$, we get the proof from the Corollary of the Fuglede-Putnam theorem [13]P279. $\square$

Corollary 4.4. *Let T be a bounded linear operator on the separable Hilbert space $\mathbb{H}$ over $\mathbb{C}$ and $T = U|T|$ is the Polar Decomposition. There is $\eta \in \mathcal{L}^1(\sigma(|T|), \mu_{|T|})$ and $\psi \in \mathcal{L}^\infty(\sigma(|T|), \mu_{|T|})$ such that $\sqrt{|\eta|} M_{z\psi}$ is the unitary equivalent representation of T on $\mathcal{L}^2(\sigma(|T|), \mu_{|T|})$, and $\sqrt{|\eta|} M_\psi$ is the unitary equivalent representation of the partial isometry U .*

Proof. For $a \in \rho(T)$, $S = T - a$ is an invertible bounded linear operator on $\mathbb{H}$, where $\rho(T) = \mathbb{C} \setminus \sigma(T)$. Because $F_{xx^*}^{\mathbb{H}}$ is a unitary operator induced by $\sqrt{|\phi|} \in \mathcal{L}^2(\sigma(|S|), \mu_{|S|})$, we get $TT^* = F_{xx^*}^{\mathbb{H}}|T|^2(F_{xx^*}^{\mathbb{H}})^*$. With [23]P54 and a similarly construction in (2) of theorem 4.1, we get that $F_{xx^*}^{\mathbb{H}}$ is a unitary operator induced by $\sqrt{|\eta|}$ on $\mathbb{H}$, where $\sqrt{|\eta|} \in \mathcal{L}^2(\sigma(|T|), \mu_{|T|})$. By the Polar Decomposition $T = U|T|$, we get that $T^*T = |T|^2$ and $TT^* = U|T|^2U^*$. Also by the proof of Theorem 4.3 we get the conclusion. $\square$

So if some chaos or dynamics or other properties of operators on Hilbert spaces only depending norm which is compatible with the inner product, then this chaos or dynamics or other properties only depending the corresponding properties of elements of $\{M_\phi : \phi \in \mathcal{L}^\infty(\sigma(|T|), \mu_{|T|})\}$, keeping the unitary equivalent representation in mind. Therefor the unitary equivalent representation is a powerful facilities, it benefit to realize the importance to deal with this problems. In the last we give some properties of normal operator through the unitary equivalent representation.

Corollary 4.5. *Let T be a bounded linear operator on the separable Hilbert space $\mathbb{H}$ over $\mathbb{C}$ and $TT^* = T^*T$. There is $\phi \in \mathcal{L}^\infty(\sigma(|T|), \mu_{|T|})$ such that $M_{z\phi}$ is the unitary equivalent representation of T on $\mathcal{L}^2(\sigma(|T|), \mu_{|T|})$. That is $T \in \mathcal{A}'(|T|)$, where $\mathcal{A}'(|T|) = \{A \in \mathcal{B}(\mathbb{H}) : AB = BA \text{ for every } B \in \mathcal{A}(|T|)\}$.*

Corollary 4.6. *Let T be a bounded linear operator on the separable Hilbert space $\mathbb{H}$ over $\mathbb{C}$, such that T is normal if and only if T is unitary equivalent to $M_{z\phi}$ on $\mathcal{L}^2(\sigma(|T|), \mu_{|T|})$, and if and only if $T \in \mathcal{A}'(|T|)$.*

REFERENCES

1. C. Kitai, *Invariant closed sets for linear operators*, Ph. D. thesis, University of Toronto, Toronto, 1982.
2. D. Stockman, *Li-Yorke Chaos in Models with Backward Dynamics*, (2012), available at <http://sites.udel.edu/econseminar/files/2012/03/lyc-backward.pdf>

3. E. M. Alfsen, *On the Dirichlet problem of the Choquet boundary*, Acta Math. **120** (1968), 149-159.
4. F. Bayart and E. Matheron, *Dynamics of Linear Operators*, Cambridge University Press 2009.
5. G. Godefroy, *Renorming of Banach spaces*, In Handbook of the Geometry of Banach Spaces, North Holland, **1** (2003), 781-835.
6. Hua Loo-kang, *On the automorphisms of a sfield*, Proc. Nat. Acad. Sci. U. S. A. 1949, 35-386-389.
7. Hou Bingzhe and Luo Lvlin, *Li-Yorke chaos for invertible mappings on noncompact spaces*, Turk. J. Math. **40** (2016), 411-416.
8. Ichiro Fujimoto, *A Gelfand representation for C^* -algebras*, Pacific J. Math. **39** (1971), 1-11.
9. Ichiro Fujimoto, *A Gelfand-Naimark Theorem FOR C^* -Algebras*, Pacific Journal of Mathematics **184(1)** (1998).
10. Izrail M. Gelfand and Mark A. Naimark, *On the embedding of normed linear rings into the ring of operators in Hilbert space*, Mat. Sbornik **12** (1943), 197-213.
11. J. H. Shapiro, *Notes on the dynamics of linear operators*, Available at the author's web page 2001.
12. J. M. G. Fell, *The structure of algebras of operator fields*, Acta Math. **106** (1961), 233-280.
13. John B. Conway, *A Course in Functional Analysis*, Second Edition, Springer-Verlag New York 1990.
14. John B. Conway, *A Course in Operator Theory*, American Mathematical Society 2000.
15. K. -G. Grosse-Erdmann, *Universal families and hypercyclic vectors*, Bull. Amer. Math. Soc. **36(3)** (1999), 345-381.
16. K. -G. Grosse-Erdmann and A. Peris Manguillot, *Linear Chaos*, Springer, London, 2011
17. Luo Lvlin and Hou Bingzhe, *Li-Yorke chaos for invertible mappings on compact metric spaces*, Arch. Math. 2016, DOI 10.1007/s00013-016-0972-5.
18. Luo Lvlin and Hou Bingzhe, *Some remarks on distributional chaos for bounded linear operators*, Turk. J. Math. **39** (2015), 251-258.
19. M. Takesaki, *A duality in the representation theory of C^* -algebras*, Ann. of Math. **85** (1967), 370-382.
20. Paul R. Halmos, *Measure Theory*, Springer-Verlag New York 1974.
21. P. Kruszyński and S.L. Woronowicz, *A non-commutative Gelfand-Naimark theorem*, J. Operator Theory **8** (1982), 361-389.
22. Robert S. Doran, *C^* -algebras: 1943-1993 (A fifty year celebration)*, (ed.) vol. 120, Contemporary Mathematics, American Mathematical Society 1994.
23. William Arveson, *A Short Course on Spectral Theory*, Springer Science+Business Media, LLC, 2002.
24. Organizers: George Elliott (University of Toronto), Thierry Giordano (University of Ottawa), David Kerr (Texas A&M University), Andrew Toms (Purdue University), <http://www.birs.ca/events/2014/5-day-workshops/14w5161>
25. Organizers: Gary Birkenmeier (UL-Lafayette), Daniel Davis (UL-Lafayette), P. W. Ng (UL-Lafayette), Leonel Robert (UL-Lafayette), Mark Tomforde (Houston), <http://www.ucs.louisiana.edu/~pwn1677/2015/2015GA.html>

SCHOOL OF MATHEMATICS AND STATISTICS, XIDIAN UNIVERSITY, 710071, XI'AN, P. R. CHINA

E-mail address: luoll12@mails.jlu.edu.cn